



ORIGINALS

Bilingual edition english/spanish

## Cysteamine ophthalmic hydrogel for the treatment of ocular cystinosis

### Hidrogel oftálmico de cisteamina para el tratamiento de la cistinosis ocular

Anxo Fernández-Ferreiro<sup>1,2,3</sup>, Andrea Luaces-Rodríguez<sup>2,3</sup>, Victoria Díaz-Tomé<sup>3</sup>, María Gil-Martínez<sup>4</sup>, María Teresa Rodríguez Ares<sup>4</sup>, Rosario Touriño Peralba<sup>4</sup>, José Blanco-Méndez<sup>3</sup>, Miguel González-Barcia<sup>1,2,3</sup>, Francisco Javier Otero-Espinar<sup>1</sup>, María Jesús Lamas<sup>2,3</sup>.

<sup>1</sup>Pharmacy Service, Xerencia Xestión Integrada Santiago de Compostela (SERGAS). <sup>2</sup>Clinical Pharmacology Group, Instituto de Investigación Sanitaria, Santiago de Compostela (IDIS-ISCIH). <sup>3</sup>Pharmacology, Pharmacy and Pharmaceutical Technology Department, Faculty of Pharmacy, University of Santiago of Compostela (USC). <sup>4</sup>Ophthalmology Service, Pharmacy Service, Xerencia Santiago de Compostela (SERGAS). Spain.

**Author of correspondence**

María Jesus Lamas

Correo electrónico:  
maria.jesus.lamas.diaz@sergas.es

Francisco Javier Otero Espinar

Correo electrónico:  
francisco.otero@usc.es

Recibido el 23 de junio de 2017;  
aceptado el 8 de agosto de 2017.

DOI: 10.7399/fh.10834

### Abstract

Ocular cystinosis is a rare disease characterised by the deposition on the corneal surface of cystine crystals that hinder the eyesight of patients. Oral cysteamine is administered as cysteamine bitartrate; however, due to the lack of corneal vascularization it does not reach the cornea and must be administered by the topical ocular route. The aim of the present study was to determine the stability of an ophthalmic hydrogel of cysteamine under different preservation conditions during a follow-up of 30 days. This hydrogel could be prepared in hospital pharmacy services.

Several physical and chemical parameters were assessed: osmolality, pH, and cysteamine concentration, which was measured using an ultra-performance liquid chromatography-tandem mass spectrometer (UPLC-MS/MS) method. Descriptive tests were also performed, such as transparency measurements and microbiological assays in order to verify sterility.

The results show that cysteamine hydrogel is stable over 30 days and should be preserved under refrigeration.

### Resumen

La cistinosis ocular es una enfermedad rara que se caracteriza por el depósito de cristales de cistina a nivel corneal, los cuales dificultan la visión de los pacientes. La cisteamina oral se administra en forma de cisteamina, pero esta no alcanza la córnea debido a la falta de vascularización corneal, por lo que es necesaria la aplicación tópica ocular. El objetivo del presente trabajo es determinar la estabilidad de un hidrogel oftálmico de cisteamina, potencialmente formulable en servicios de farmacia hospitalaria, conservado este bajo diferentes condiciones de almacenamiento durante un periodo de 30 días.

Los parámetros físicos y químicos evaluados han sido la osmolalidad, el pH y la concentración de cisteamina, siendo esta última valorada mediante un método de cromatografía líquida de ultra alta presión, empleando un detector de masas en tándem (UPLC-MS/MS). Los ensayos descriptivos se han basado en la medición de la transparencia y los ensayos microbiológicos en la realización de pruebas de esterilidad.

Los resultados obtenidos permiten concluir que el hidrogel de cisteamina es estable durante un periodo de 30 días, recomendándose que su conservación sea en nevera.

### KEYWORDS

Cysteamine; Ellman's reagent; Hyaluronic acid; Mass spectrometry; Ocular cystinosis; Ophthalmic hydrogel; Stability.

### PALABRAS CLAVE

Ácido hialurónico; Cisteamina; Cistinosis; Ocular; Estabilidad; Espectrometría de masas; Hidrogel oftálmico; Reactivo de Ellman.



Los artículos publicados en esta revista se distribuyen con la licencia  
Articles published in this journal are licensed with a  
Creative Commons Attribution 4.0  
<https://creativecommons.org/licenses/by-nc-nd/4.0/>  
La revista Farmacia no cobra tasas por el envío de trabajos,  
ni tampoco por la publicación de sus artículos.

## Introduction

Cystinosis is a rare lysosomal storage disorder that follows an autosomal recessive inheritance pattern<sup>1</sup>. This metabolic disorder is characterized by the accumulation of the amino acid cystine in lysosomes due to defective cystine transportation from the interior to the exterior of the lysosome<sup>2,3</sup>. Cystine has low solubility in water, leading to the formation of intralysosomal crystals and damage to various tissues and organs, including the cornea. As described by Burki in the 1940s, the ocular manifestations of the disease are due to the accumulation of cystine crystals in the ocular surface<sup>4</sup>. These crystals can be observed with a slit lamp and are a pathognomonic sign of cystinosis. They begin to form during infancy and from 16 months of age onward they can be observed through a slit lamp. Patients are initially asymptomatic. Due to the accumulation of corneal cystine crystals over time, ocular symptoms do not appear until approximately 10 years of age<sup>5</sup>.

The specific treatment of cystinosis is cysteamine, also called mercaptamine or 2-aminoethanethiol, which is an aminothiols with chemical formula  $\text{HSCH}_2\text{CH}_2\text{NH}_2$ <sup>6</sup>. Cysteamine was introduced as a possible therapeutic agent for cystinosis in 1976 and remains the only available treatment<sup>7</sup>. Although cysteamine does not cure cystinosis, it has revolutionized patient management and prognosis. It has been shown to slow disease progression and can reduce the amount of intracellular cystine by more than 90%. Cysteamine therapy should be started as soon as the diagnosis is made and should be continued for the lifetime of the patient. Patients with poor adherence to treatment or begin it later do not achieve such beneficial outcomes<sup>8</sup>.

Oral cysteamine is administered in the form of cysteamine bitartrate, but does not reach the cornea due to the lack of corneal vascularization. Thus, a topical ocular application was developed, whose safety and effectiveness had already been demonstrated in the 1980s<sup>9-11</sup>. Currently, there are two available ophthalmic formulations of cysteamine hydrochloride: Cystaran® (Sigma Tau Pharmaceuticals Inc.), an FDA-approved medication, which must be instilled from 6 to 12 times a day<sup>12</sup>; and Cystadrops® (Orphan Europe, Paris, France), which has a higher viscosity and increased ocular permanence<sup>13,14</sup>. Cystadrops is currently in Phase III trials; however, the European Medicines Agency has recently allowed it be marketed as an orphan drug to facilitate access<sup>15</sup>.

Access to foreign and/or orphan drugs can sometimes be delayed by the obligatory procedures and approvals required for their use. In addition, the sometimes exorbitant price of these drugs can hamper access<sup>16</sup>. In order to facilitate the treatment of ocular cystinosis, cysteamine eye drops as a compounded formulation are commonly prepared in hospital pharmacy services.

Two major problems are associated with these formulations. Firstly, cysteamine eye drops must be instilled every hour while the patient is awake to reduce the amount of corneal crystals. To optimise the formulation and avoid these difficult dosage schedules, our group developed a bioadhesive cysteamine hydrogel with high ocular permanence, which could be prepared by hospital pharmacy services<sup>17,18</sup>. Secondly, there is a lack of studies on the stability of cysteamine formulations. The analysis of compounds with thiol groups has always proved difficult, owing to their susceptibility to oxidation and the lack of a structural chromophore needed for their detection<sup>19,20</sup>. Furthermore, the low molecular weight of cysteamine (MW = 77.15 g/mol) hinders its direct detection by mass detectors. Thus, the methods used to determine these types of compounds usually derivatize the cysteamine molecule before quantification<sup>21</sup>.

The objective of this article was to determine the stability of a bioadhesive ophthalmic cysteamine hydrogel under different storage conditions.

## Methods

### Preparation of 0.55% ophthalmic cysteamine hydrogel

The preparation of the hydrogel is performed in 2 stages. Firstly, a sufficient quantity of cysteamine (BioXtra, Sigma-Aldrich) is gradually added to Balanced Salt Solution Alcon® and magnetically stirred over a period of 5 minutes to achieve a concentration of 0.55%. While continuing to stir,

hyaluronic acid (Acofarma®) is then added to achieve a final concentration of 0.4%.

Secondly, the resulting hydrogel is vacuum filtered using a 0.22- $\mu\text{m}$  membrane filter (Stericup® Merck Millipore Express™ PLUS 0.22  $\mu\text{m}$ ) and poured into 15-ml type-1 amber glass containers, adding 10 mL of hydrogel to each container. The remaining volume is filled with nitrogen gas, and the containers are closed. The entire process is performed under aseptic conditions using a horizontal laminar flow hood.

### Preservation conditions and study variables

The formulas were divided into 2 batches: those without preservatives and those with preservatives. The latter batches were prepared by adding 0.01% Ethylene Diamine Tetraacetic Acid (EDTA) while dissolving the cysteamine. Half of the batches with and without preservatives were stored for 30 days at 22°C (room temperature) and the other batches were stored at 4°C (refrigerated). In the rest of this article, these formulations are referred to as HA (room temperature without EDTA), HAE (room temperature with EDTA), HN (refrigerated without EDTA), and HNE (refrigerated with EDTA).

Osmolality, pH, and cysteamine concentrations were assessed. Descriptive tests were based on transparency measurements, and microbiological tests were based on sterility testing.

All samples were allowed to reach and stay at room temperature for a minimum of 30 minutes to avoid measurement errors due to temperature variations. All tests were performed in triplicate and were conducted on days 0, 7, 14, and 30 after the preparation of the hydrogels.

### Descriptive tests

The transparency of the samples was determined by measuring transmittance in the visible light range (380 nm - 780 nm) using a UV-VIS spectrophotometer (model 8452 Diode Array Spectrophotometer, Hewlett Packard). A blank was made with distilled water, the different formulations were placed in quartz cuvettes, and then transmittance was measured, thus obtaining a graph representing the percentage of transmitted light as a function of wavelength.

### Physicochemical tests

**Determination of osmolality and pH:** Osmolality was measured using a vapour pressure osmometer (VAPRO 5520). 10  $\mu\text{L}$  of each formulation was deposited on a disk of Whatman filter paper on the chamber. pH was determined using a Crison microPH2001® pH-metre.

**Quantification of cysteamine:** A saturated solution of Ellman's reagent (5,5'-dithiobis(2-nitrobenzoic acid)) was prepared as a derivatizing agent<sup>22</sup> (Figure 1), by dissolving 29.6 mg of the powder in 10 mL of 0.018 M aqueous NaOH. Subsequently, the solution was filtered through a 0.45  $\mu\text{m}$  filter.

To quantify cysteamine, a 1:1000 dilution of the formulation was prepared. 100  $\mu\text{L}$  of Ellman's reagent and 100  $\mu\text{L}$  of purified water were added to this diluted sample. The resulting solution was analysed using an ultra-performance liquid chromatography-tandem mass spectrometer (UPLC-MS/MS) method. Measurements were taken using an Acquity UPLC® H-Class system (Waters® Milford, Massachusetts) coupled to a Xevo TQD mass spectrometer (Waters®). Data were collected using Mass lynx v4.1 software and processed using Target lynx™ Application Manager chromatographic software. Chromatographic separation was conducted at 40°C using an Acquity BEH C<sub>18</sub> column (2.1 mm x 50 mm; particle size 1.7  $\mu\text{m}$ ) (Waters®). The mobile phase solvents used were a 0.1% formic acid solution in water (MilliQ®) (Phase A) and acetonitrile (Phase B). A gradient with constant flow rate of 0.4 mL/min was used. The gradient was started at 100% phase A, changing linearly to a 40% A – 60% B composition at 2.2 minutes, maintaining the composition until the 2.60-minute mark, and then returned to initial conditions at 3 minutes. The autosampler was set to 10°C and 10  $\mu\text{L}$  of each sample was injected. Total run time was 3 minutes, which included equilibration of the chromatographic system prior to sample injection. Mass spectrometry data were obtained using the multiple reaction monitoring (MRM) mode through positive electrospray ionization. Quantification was achieved by means of the transitions of




Figure 1. Cysteamine derivatization using Ellman's reagent.

the precursor ion at  $m/z$  313 and the 196.85 fragment ion using a desolvation gas flow rate of 1.1 L/h, cone gas flow rate of 80 L/h, and a capillary voltage of 3.2 kV. Desolvation and source temperatures were 600°C and 146°C, respectively.

### Microbiological tests

Each of the hydrogels was analysed on the aforementioned days to determine microbiological stability. 1 mL of each of the hydrogels was added to plates containing blood agar, sabouraud agar, and fluid thioglycolate medium. The samples were cultured at 37°C for 48 hours, 15 days, and 10 days, respectively. At the end of each incubation period, the samples were inspected for any signs of microbiological growth.

### Allowed variation range and statistical analysis

The Pharmaceutical Codex was used to establish the expiry date of the formulation, which was set<sup>23</sup> when there was a 10% reduction of active ingredients compared to the initial concentration. Changes in pH and osmolality were considered unacceptable if their values exceeded the acceptance criteria for ophthalmic applications. Microbiological stability was considered acceptable providing no microbial growth was detected in the cultured samples. Finally, the product was considered unacceptable in the absence of complete transparency on descriptive tests.

The results of the different preservation conditions were compared by multivariate analysis of variance using Graph Pad Prism® v.5.0b software.

## Results

### Descriptive and physicochemical tests

All the formulations were completely transparent and no decrease in transparency was observed over the study period. No signal was observed in the visible range, demonstrating the transparency of the sample (see figure 2).

Figure 3 shows variations in osmolality of the cysteamine hydrogel under all four preservation conditions over time. Osmolality values of all formulations remained between 90% and 100% of the initial values over the study period. Under the different study conditions, no statistically significant differences were observed between the formulations, although those containing EDTA showed slightly higher values ( $427 \pm 8.96$  mOsm/Kg vs  $410 \pm 9.48$  mOsm/Kg).

However, as shown in figure 4, neither the addition of EDTA to the hydrogel nor storage temperature influenced the pH values of the hydrogel over the study period. Under all the conditions tested, no statistically significant differences were observed between the initial and final pH, except for a slight decrease in pH in the EDTA formulations (6.29 vs 6.44).

### Concentration of cysteamine

A narrow, symmetrical, and well-defined chromatographic peak was obtained with an elution time of 0.33 minutes. Figure 5 shows an example chromatogram of the derivatised cysteamine obtained using the UPLC-MS/MS method.





Figure 2. Graph obtained by determining the transparency of one of the formulations, showing negligible absorbance in the visible light spectrum (380 nm - 780 nm).




**Figure 3.** Percentage change in osmolality (mOsm/Kg) (mean and standard deviation) of cysteamine hydrogel over 30 days under the four different storage conditions. HA, room temperature without EDTA; HAE, room temperature with EDTA; HN, refrigerated without EDTA; HNE, refrigerated with EDTA.

The UPLC-MS/MS determination method employed is highly specific, because it combines the efficiency of chromatographic separation and the high selectivity of a tandem mass detector to select the chemical structure to be determined. Using this method, the derivatised product was separated from any compounds that may have formed from cysteamine degradation.


Figure 6 shows variations in cysteamine concentrations over time under the four different storage conditions. Cysteamine concentrations did not fall below 90% at any point during the study period. Nevertheless, it should be noted that a wide range of concentration values was observed at different time points during the storage period. Some samples had percentages of more than 100% of the initial concentration. This variability was caused by the high viscosity of hydrogels, which makes it difficult to obtain reproducible volumetric samples by aspiration.

### Microbiological stability

Adequate storage of samples was maintained under all study conditions, and no microbial growth was observed in any of the hydrogels during the storage period.



**Figure 4.** Change in pH (mean and standard deviation) of cysteamine hydrogel over 30 days under the four different storage conditions. HA, room temperature without EDTA; HAE, room temperature with EDTA; HN, refrigerated without EDTA; HNE, refrigerated with EDTA.




**Figure 5.** Example of chromatogram obtained for cysteamine derivatised with Ellman's reagent, using the UPLC MS/MS method.

### Discussion

Difficult dosage schedules are the major challenge to the use of ophthalmic compounded formulations of cysteamine hydrochloride, because they require hourly instillations to reduce the amount of corneal crystals. A preclinical study has shown that the biopermanence of the hydrogel under study is similar to that of Cystadrops®<sup>18</sup>. Quantitative PET biopermanence studies have shown that cysteamine hydrogel with hyaluronic acid has a 60-minute half-life, which is much higher than the 18-minute half-life of cysteamine eye drops typically prepared by hospital pharmacy services. In addition, more cysteamine reaches the stroma after administration of this hydrogel than reaches it with eye drops, and there are statistically significant differences in transcorneal permeation values between the two media. This hydrogel formulation achieves a controlled release of cysteamine over time, and can be prepared by pharmacy services for patients with ocular cystinosis<sup>17</sup>. Two preparation methods are presented in the supplementary materials.

Stability studies are a relevant technical and economic challenge for hospital pharmacy services and are becoming more frequent to guarantee the



**Figure 6.** Percentage of change in cysteamine concentrations over 30 days. HA, room temperature without EDTA; HAE, room temperature with EDTA; HN, refrigerated without EDTA; HNE, refrigerated with EDTA.

quality of the prepared drugs<sup>24</sup>. The present study investigated the stability of cysteamine hydrogel with hyaluronic acid, and showed that its properties were unchanged in a variety of storage conditions over the study period. Other authors have shown that at room temperature cysteamine oxidises into its cystamine dimer, which is ineffective for the removal of corneal cystine crystals<sup>25</sup>. For this reason, nitrogen was used to remove environmental oxygen before sealing the containers.

The choice of EDTA as a preservative was based on previous publications, which have shown it to be the most suitable preservative for use in cysteamine formulations. The lowest possible concentration of EDTA was chosen to minimize potential toxicity on the corneal epithelium<sup>20, 26-28</sup>. Other preservatives, such as benzalkonium chloride, were ruled out after an unfavourable benefit-risk assessment by ophthalmologists. This decision was based on their potential epithelial toxicity, which would be higher during the chronic use of the hydrogel under study<sup>29</sup>.

According to the United States Pharmacopoeia, the stability of a compounded formulation is defined as the amount of time during which a product maintains, within very specific limits, the properties and characteristics that it possessed at the time of manufacture, throughout storage, and during use<sup>30</sup>. Stability studies determine how the quality of a drug varies over time under the influence of a number of factors, and use this information to provide recommendations on its expiration date and storage conditions. Over the study period, the pH and osmolality of the hydrogels remained practically constant, with no statistically significant differences ( $<0.001$ ) between the initial and final values under all storage conditions. The addition of EDTA to the formulation led to a slight increase in osmolality and a decrease in pH without affecting the stability of cysteamine. All the measures applied showed that transparency was 100%. Cysteamine hydrogel with hyaluronic acid maintained its properties for 30 days after preparation. However, because the addition of EDTA did not improve stability and the use of benzalkonium chloride as a preservative was discarded, it is recommended that the hydrogel should be stored in a refrigerator to prevent microbiological growth and as an alternative method to sealing the containers with nitrogen<sup>31</sup>.

The properties of cysteamine hydrogel have been described in previous studies and the results of this stability study show that the use of cysteamine

hydrogel may increase therapeutic benefit in patients with ocular cystinosis. In addition, this formulation may be an effective alternative to those not marketed in Spain, but which are available for importation. The cost of imported formulations is €37,728/y/patient, whereas the estimated cost of producing the formulation is €1,080/y/patient.

The authors conducted a survey using the mailing list of the Spanish Society of Hospital Pharmacy (SEFH) and estimated that there are currently 39 patients under treatment with ocular topical cysteamine in Spain. Thus, the cost of this drug acquired through the Access to Medicines not Authorized in Spain process would be €1,471,392/y. The use of cysteamine hydrogel prepared in hospital pharmacy services would provide patients with better access to treatment and achieve significant savings for the Spanish National Health System.

## Acknowledgements

- *Funding for staff hiring:* A.F-F, Carlos III Health Institute (ISCIII) CM15/00188
- *Funding for research projects:* Mutua Madrileña Foundation and the Spanish Foundation of Hospital Pharmacy.
- *Special thanks to:* All SEFH members who responded to the survey on the use of ophthalmic cysteamine in Spain. This survey was conducted using the mailing list of the SEFH.

## Contribution to scientific literature

Cysteamine eye drops are commonly prepared as a compounded formulation in hospital pharmacy services. Two problems are associated with these formulations: their low permanence at the ocular level and the lack of studies on their stability. This article is the first study on the stability of a cysteamine hydrogel with high ocular permanence, which could be prepared by hospital pharmacy services, and may therefore represent a great advance in the treatment of ocular cystinosis.

## Bibliography

1. Gahl WA, Thoene JG, Schneider JA. Cystinosis. *N Engl J Med.* 2002;347(2):111-21. doi:10.1056/NEJMra020552
2. Linkage of the gene for cystinosis to markers on the short arm of chromosome 17. The Cystinosis Collaborative Research Group. *Nat Genet.* 1995;10(2):246-8. doi:10.1038/ng0695246
3. Town M, Jean G, Cherqui S, Attard M, Forestier L, Whitmore SA, et al. A novel gene encoding an integral membrane protein is mutated in nephropathic cystinosis. *Nat Genet.* 1998;18(4):319-24. doi:10.1038/ng0498319
4. Burki E. Ueber die Cystinkrankheit im Kleinkindesalter unter besonderer Berücksichtigung des Augenbefundes [About the Cystinosis in infancy with special reference to eye findings]. *Ophthalmologica.* 1941;101:331-42.
5. Gahl WA, Kuehl EM, Iwata F, Lindblad A, Kaiser-Kupfer MI. Corneal crystals in nephropathic cystinosis: natural history and treatment with cysteamine eyedrops. *Mol Genet Metab.* 2000;71(1-2):100-20. doi:10.1006/mgme.2000.3062
6. Pubchem. Cysteamine (C2H7NS). [Accessed June 22, 2017] Disponible en: <https://pubchem.ncbi.nlm.nih.gov/compound/2-Aminoethanethiol>
7. Thoene JG, Oshima RG, Crowhall JC, Olson DL, Schneider JA. Cystinosis. Intracellular cystine depletion by aminoethiols in vitro and in vivo. *J Clin Invest.* 1976;58(1):180-9. doi:10.1172/JCI108448
8. Brodin-Sartorius A, Tête M-J, Niaudet P, Antignac C, Guest G, Ottolenghi C, et al. Cysteamine therapy delays the progression of nephropathic cystinosis in late adolescents and adults. *Kidney Int.* 2012;81(2):179-89. doi:10.1038/ki.2011.277
9. Kaiser-Kupfer MI, Caruso RC, Minkler DS, Gahl WA. Long-term ocular manifestations in nephropathic cystinosis. *Arch Ophthalmol.* 1986;104(5):706-11. doi:10.1001/archoph.1986.01050170096030
10. Cysteamine Eye Drops to Treat Corneal Crystals in Cystinosis. *Clinical Trial.* Clinicaltrials.gov [A service of the U.S. National Institutes of Health]. [Accessed May 6, 2016] Disponible en: <https://clinicaltrials.gov>
11. Kaiser-Kupfer MI, Gazzo MA, Dattiles MB, Caruso RC, Kuehl EM, Gahl WA. A randomized placebo-controlled trial of cysteamine eye drops in nephropathic cystinosis. *Arch Ophthalmol.* 1990;108(5):689-93. doi:10.1001/archoph.1990.01070070075038
12. Cystaran®. Prescribing information. [Accessed May 3, 2016] Disponible en: [www.sigmatau.com](http://www.sigmatau.com)
13. Cystadrops® 0,55% eye drops solution. Summary of product characteristics. November 2015. [Accessed September 24, 2017] Disponible en: <https://www.medicines.org.uk/emc/medicine/34065>
14. Cystadrops® 0,55% collyre en solutions. Protocole d'utilisation thérapeutique et de recueil d'informations. Autorisation temporaire d'utilisation dite de cohorte. [Accessed June 2016] Disponible en: [http://ansm.sante.fr/var/ansm\\_site/storage/original/application/f0f8036949f10b1ce8dd7a7a0ab48927.pdf](http://ansm.sante.fr/var/ansm_site/storage/original/application/f0f8036949f10b1ce8dd7a7a0ab48927.pdf)
15. Cysteamine Hydrochloride for nephropathic Cystinosis, open-label Phase III pivotal study (CYSTADROPS CHOC study). *Clinical Trials Register.* [Accessed June 18, 2016] Disponible en: [www.clinicaltrialsregister.eu](http://www.clinicaltrialsregister.eu)
16. Velásquez G. El acceso global a los medicamentos en el contexto internacional actual. *Biomédica.* 2011;31(2):162-3.
17. Luaces-Rodríguez A, Díaz-Tomé V, González-Barcia M, Silva-Rodríguez J, Herranz M, Gil-Martínez M, et al. Cysteamine polysaccharide hydrogels: study of extended ocular delivery and biopermanence time by PET imaging. *Int J Pharm.* 2017;528(1-2):714-22. doi:10.1016/j.ijpharm.2017.06.060
18. Fernández Ferreiro A, Luaces-Rodríguez A, González Barcia M, Otero Espinar FJ, Lamas MJ. Evaluación de la biopermanencia ocular in vivo de tres formulaciones oftálmicas de cisteamina clorhidrato. *Farm Hosp.* 2016;Supl1:79.
19. Guan X, Hoffman B, Dwivedi C, Matthees DP. A simultaneous liquid chromatography/mass spectrometric assay of glutathione, cysteine, homocysteine and their disulfides in biological samples. *J Pharm Biomed Anal.* 2003;31(2):251-61.
20. Lawal B. Development of a cysteamine in situ gelling system for the treatment of corneal crystals in cystinosis. [Published July 2008. Accessed February 3, 2016]. Disponible en: [https://cystinosis.org/images/research/updates/CRN\\_Research\\_UpdateB.pdf](https://cystinosis.org/images/research/updates/CRN_Research_UpdateB.pdf)
21. Qi B-L, Liu P, Wang Q-Y, Cai W-J, Yuan B-F, Feng Y-Q. Derivatization for liquid chromatography-mass spectrometry. *TrAC Trends Anal Chem.* 2014;59:121-32. doi:10.1016/j.trac.2014.03.013
22. Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys.* 1959;82(1):70-7.

23. Royal Pharmaceutical Society of Great Britain Dept of Pharmaceutical. The Pharmaceutical Codex: Principles and Practice of Pharmaceutics. Twelfth Edition. London: Pharmaceutical Press; 1994.
24. Barrueco N, Escobar Rodríguez I, García Díaz B, Gil Alegre ME, López Lunar E, Ventura Valares MG. Estabilidad de medicamentos en la práctica clínica: de la seguridad a la eficiencia. *Farm Hosp.* 2013;37(3):175-7. doi:10.7399/FH.2013.37.3.587
25. Iwata F, Kuehl EM, Reed GF, McCain LM, Gahl WA, Kaiser-Kupfer MI. A randomized clinical trial of topical cysteamine disulfide (cystamine) versus free thiol (cysteamine) in the treatment of corneal cystine crystals in cystinosis. *Mol Genet Metab.* 1998;64(4):237-42. doi:10.1006/mgme.1998.2725
26. Herreros JMA. Preparación de medicamentos y formulación magistral para oftalmología. Ediciones Díaz de Santos; 2003.
27. Fernández MA, Atienza JM, Vayo CÁ del. *Formulación en farmacia pediátrica.* Madrid. A. Madrid Vicente editores; 2011.
28. Asociación de Formelistas de Andalucía. [Accessed May 28, 2016] Disponible en: <http://www.formulistasdeandalucia.es/noticia.php?id=264>
29. Rosin LM, Bell NP. Preservative toxicity in glaucoma medication: clinical evaluation of benzalkonium chloride-free 0.5% timolol eye drops. *Clin Ophthalmol Auckl NZ.* 2013;7:2131-5. doi:10.2147/OPHT.S41358
30. Convention USP. *Farmacopea de los Estados Unidos de América, USP 37, 2014: Formulario nacional, NF 32.* United States Pharmacopeia; 2013.
31. *Practical Pharmaceutics. An International Guideline for the Preparation, Care and Use of Medicinal Products.* Yvonne Bouwman-Boer, V'lain Fenton-May, Paul Le Brun, editors. Springer 2015. [Accessed June 22, 2017]. Disponible en: <http://www.springer.com/gp/book/9783319158136>

## Appendix 1



Xestión Integrada  
 Santiago de Compostela

Xestión Integrada de Santiago de Compostela  
 University Hospital Santiago de Compostela  
 Pharmacy Service

### PHARMACEUTICAL COMPOUNDING PROTOCOL

#### A) CYSTEAMINE HYDROGEL WITH HYALURONIC ACID 0.55%

Dosage form: Ophthalm. hydrogel	Shelf life: 30 days	
Administrat. route: Topical ocular	Prot. from light: Yes	Validation date: 06/21/2017
Compounding by: Pharmacist	Storage: Refrigerated (2 - 8 °C)	Dispensable: Yes

#### Additional information

Dispose 7 days after opening.

Risk level: **MEDIUM**  
 Compounding: **Sterile**

#### Ingredients

Ingredient	Quantity
Cysteamine hydrochloride	275.00 mg
Sodium hyaluronate (powder)	40.00 mg
BSS® eye drops (Balanced Salt Solution)	q.s. 50.00 ml

#### Equipment

- 1 - Sterile plastic cup
- 1 - Analytical balance
- 1 - Magnetic stirrer
- 1 - 60-mL Luer-Lock syringe
- 1 - 0.22-µm filter (Millipore Millex® GS) or (Stericup® Merck Millipore Express™ Plus)

#### Directions

- Weigh the cysteamine and sodium hyaluronate in 100 mL sterile plastic cups.
- In a horizontal laminar flow hood, add 40 mL of BSS® to the cup containing the cysteamine. Stir in a magnetic stirrer with a previously sterilized stirring tablet until completely dissolved.
- Once ready, slowly add the hyaluronic acid, continuing to stir until completely dissolved.
- Transfer with a syringe and add BSS® to 50 mL.
- Transfer the entire content (50 mL) back to the plastic cup and continue magnetic stirring until homogeneous.
- Finally, use a syringe to transfer the 50 mL of cysteamine hydrogel and perform sterilizing filtering, storing the sterilized content in amber glass bottles. Note: Filtering of viscous substances can be expensive with conventional filters. Therefore, vacuum filtering is used as an alternative:
  - o Option 1: For small volumes use Millipore Millex® GS).
  - o Option 2: For larger volumes use Stericup® Merck Millipore Express™ PLUS 0.22 µm for vacuum filtering.
- Seal and label bottle.

**Packaging:** Amount made, 50 mL distributed into 5 sterile 15-mL glass dropper bottles each filled with 10 mL.

#### Quality Control

Test

**Particulate testing**

**Microbiological analysis**



**Xestión Integrada  
Santiago de Compostela**

Xestión Integrada de Santiago de Compostela  
**University Hospital Santiago de Compostela**  
Pharmacy Service

**PHARMACEUTICAL COMPOUNDING PROTOCOL**

**A) CYSTEAMINE HYDROGEL WITH HYALURONIC ACID 0.55%**

---

**Authors**

---

- Anxo Fernández Ferreiro (Pharmacist)
  - Miguel González Barcia (Pharmacist)
- 

**References**

---

- 1.- Andrea Luaces-Rodríguez y cols. Cysteamine polysaccharide hydrogels: study of extended ocular delivery and biopermanence time by PET imaging, *Int. J. Pharm.* In Press (2017).
- 2.- Anxo Fernández Ferreiro y cols. Evaluación de la biopermanencia ocular in vivo de tres formulaciones oftálmicas de cisteamina clorhidrato., *Farm. Hosp.* 61 SEFH Congress (2016).
- 3.- Manuela Atienza Fernández. *Formulación en Farmacia Pediátrica*. 2005 3rd edition - ISBN: 84- 689- 2529- 2. 4. - M.I. Kaiser-Kupfer, M.A. Gazzo, M.B. Datiles, R.C. Caruso, E.M. Kuehl, W.A. Gahl, A randomized placebo-controlled trial of cysteamine eye drops in nephropathic cystinosis, *Arch. Ophthalmol. Chic.* III 1960. 108- 1990- 689- 693.

## Appendix 2



Xestión Integrada  
 Santiago de Compostela

Xestión Integrada de Santiago de Compostela  
 University Hospital Santiago de Compostela  
 Pharmacy Service

### PHARMACEUTICAL COMPOUNDING PROTOCOL

#### B) CYSTEAMINE HYDROGEL WITH HYALURONIC ACID 0.55%

Dosage form: Ophthalm. hydrogel	Shelf life: 30 days	Validation date: 06/21/2017
Administrat. route: Topical ocular	Prot. from light: Yes	Dispensable: Yes
Compounding by: Pharmacist	Storage: Refrigerated (2 - 8 °C)	
<b>Additional information</b> Dispose 7 days after opening.		Risk level: <b>MEDIUM</b> Compounding: <b>Sterile</b>

#### Ingredients

Ingredient	Quantity
Cysteamine hydrochloride	275.00 mg
BSS® eye drops (Balanced Salt Solution)	5.00 ml
Sterile artificial tear drops with hyaluronic acid 0.4% Acuoral®	q.s. 50.00 ml

#### Equipment

- 1 - Sterile plastic cup
- 1 - Analytic balance
- 1 - Magnetic stirrer
- 1 - 60 mL Luer®-Lock syringe
- 1 - 0.22-µm filter (Millipore Millex® GS)

#### Directions

- Weigh the cysteamine in a sterile plastic cup.
- In a horizontal laminar flow hood, add 5 mL of BSS® to the cup containing cysteamine. Mix in a magnetic stirrer with a previously sterilized stirring tablet, until completely dissolved.
- Transfer the cysteamine solution to a syringe and perform sterilizing filtering (Millipore Millex® GS), then transfer the filtered content to a 60-mL syringe.
- Next, add artificial tear drops to complete 50 mL.
- Homogenize by moving the syringe and then distribute into amber glass bottles.
- Seal and label bottles.

**Packaging:** Quantity to manufacture, 50 mL distributed in 5 sterile 15-mL glass dropper bottles each filled with 10 mL

#### Quality Control

##### Test

##### Particulate testing

##### Microbiological analysis

#### Authors

- Anxo Fernández Ferreiro (Pharmacist)
- Miguel González Barcia (Pharmacist)

#### References

- 1.- Andrea Luaces-Rodríguez y cols. Cysteamine polysaccharide hydrogels: study of extended ocular delivery and biopermanence time by PET imaging. Int. J. Pharm. In Press (2017).
- 2.- Anxo Fernández Ferreiro y cols. Evaluación de la biopermanencia ocular in vivo de tres formulaciones oftálmicas de cisteamina clorhidrato., Farm. Hosp. 61 SEFH Congress (2016).



SERVIZO  
GALEGO  
de SAÚDE

Xestión Integrada  
Santiago de Compostela

Xestión Integrada de Santiago de Compostela  
University Hospital Santiago de Compostela  
Pharmacy Service

**PHARMACEUTICAL COMPOUNDING PROTOCOL**

**B) CYSTEAMINE HYDROGEL WITH HYALURONIC ACID 0.55%**

3. - Manuela Atienza Fernández. Formulación en Farmacia Pediátrica. 2005 3rd edition - ISBN: 84- 689- 2529- 2. 4. - M.I. Kaiser-Kupfer, M.A. Gazzo, M.B. Datiles, R.C. Caruso, E.M. Kuehl, W.A. Gahl, A randomized placebo-controlled trial of cysteamine eye drops in nephropathic cystinosis, Arch. Ophthalmol, Chic. Ill 1960. 108- 1990- 689- 693.